

BRIEF REPORT

WILEY

Community norms for the Eating Disorder Examination Questionnaire among cisgender gay men

Jason M. Nagata¹ | Matthew R. Capriotti^{2,3,4} | Stuart B. Murray⁵ |
 Emilio J. Compte^{5,6,7} | Scott Griffiths⁸ | Kirsten Bibbins-Domingo^{3,9} |
 Juno Obedin-Maliver^{4,10} | Annesa Flentje^{4,5,11} | Micah E. Lubensky^{4,12} |
 Mitchell R. Lunn^{4,12}

¹Department of Pediatrics, University of California, San Francisco, California, USA

²Department of Psychology, San José State University, San Jose, California, USA

³Department of Medicine, University of California, San Francisco, California, USA

⁴The PRIDE Study/PRIDEnet, Stanford University School of Medicine, Stanford, California, USA

⁵Department of Psychiatry, University of California, San Francisco, California, USA

⁶School of Human and Behavioral Sciences, Favaloro University, Buenos Aires, Argentina

⁷DBT-Eating Disorders Team, Fundación Foro, Buenos Aires, Argentina

⁸Melbourne School of Psychological Sciences, University of Melbourne, Victoria, Australia

⁹Department of Epidemiology and Biostatistics, University of California, San Francisco, California, USA

¹⁰Department of Obstetrics and Gynecology, Stanford University School of Medicine, Stanford, California, USA

¹¹Department of Community Health Systems, University of California, San Francisco, California, USA

¹²Division of Nephrology, Department of Medicine, Stanford University School of Medicine, Stanford, California, USA

Correspondence

Jason M. Nagata, M.D., M.Sc., 550 16th Street, 4th Floor, Box 0110, San Francisco, CA 94158.

Email: jasonmnagata@gmail.com

Funding information

American Heart Association, Grant/Award Number: CDA34760281; Eunice Kennedy Shriver National Institute of Child Health and Human Development, Grant/Award Number: K12 HD000850; National Health and Medical Research Council, Grant/Award Number: 1121538; National Institute of Diabetes and Digestive and Kidney Diseases, Grant/Award Number: K12DK111028; National Institute of Mental Health, Grant/Award Number: K23 MH115184; Patient-Centered Outcomes Research Institute, Grant/Award Number: PPRN-1501-26848; National Institute on Drug Abuse, Grant/Award Number: K23DA039800; American Pediatric Society; American Academy of Pediatrics

Abstract

Objective: Prior norms of the Eating Disorders Examination Questionnaire (EDE-Q) among men have not considered sexual orientation. This study's objective was to assess EDE-Q community norms among cisgender gay men.

Method: Participants were 978 self-identified cisgender gay men from The PRIDE Study recruited in 2018.

Results: We present mean scores and standard deviations for the EDE-Q among cisgender gay men ages 18–82. Among cisgender gay men, 4.0% scored in the clinically significant range on the global score, 5.7% on the restraint, 2.1% on the eating concern, 10.5% on the weight concern, and 21.4% on the shape concern subscales of the EDE-Q. The global score as well as weight and shape concerns in a young adult subsample (18–26 years) from The PRIDE Study were higher than previously reported norms in young men (Lavender, 2010). Participants reported any occurrence ($\geq 1/28$ days) of dietary restraint (19.8%), objective binge episodes (10.9%), excessive exercise (10.1%), laxative misuse (1.1%), and self-induced vomiting (0.6%). Binge eating, excessive exercise, and self-induced vomiting in The PRIDE Study subsample were lower than previously reported in young men.

Discussion: We provide EDE-Q norms among cisgender gay men, which should aid clinicians and researchers to interpret the EDE-Q scores of cisgender gay men.

KEYWORDS

body image, disordered eating, eating disorder, gay, homosexuality, norms, sexual minority

1 | INTRODUCTION

Most empirical eating disorder (ED) research is skewed towards populations of women with less than 1% of all published ED research relating specifically to men (Murray et al., 2017). Despite this skew, emerging evidence suggests that EDs are increasingly prevalent in men (Mitchison & Mond, 2015) and have disease burden and medical risk comparable with that of women (Murray et al., 2017). Accordingly, an increasing research impetus has been oriented towards EDs in men, both in terms of improving detection and optimizing treatment. However, men may have unique ED-related concerns, such as muscularity versus thinness-oriented body image ideals (Darcy, Hardy, Lock, Hill, & Peebles, 2013; Murray et al., 2017). As such, revised normative data are required to ensure appropriate measure interpretation in men (Lavender, De Young, & Anderson, 2010).

Although revised norms for commonly used measures of ED symptomatology—such as the Eating Disorder Examination Questionnaire (EDE-Q; Fairburn & Beglin, 2008)—have been put forth for men in clinical and community (i.e., non-clinical) settings (Hilbert, de Zwaan, & Braehler, 2012; Lavender et al., 2010; Quick & Byrd-Bredbenner, 2013; Reas, Øverås, & Rø, 2012; Smith et al., 2017), these revised norms have not considered the respondents' sexual orientation. Although evidence suggests an association between gay sexual orientation and elevated disordered eating in men (Austin et al., 2009; Calzo, Austin, & Micali, 2018; Matthews-Ewald, Zullig, & Ward, 2014), there is limited evidence that gay sexual orientation is a direct risk factor in the development of EDs among men (Siever, 1994). In one study, gay adolescent boys in high school were more likely to perceive themselves as overweight when they were normal weight (Hadland, Austin, Goodenow, & Calzo, 2014). Gay undergraduate men had a higher incidence of body dissatisfaction and a drive for thinness than their heterosexual counterparts (Carper, Negy, & Tantleff-Dunn, 2010). Sexual orientation-specific community norms may contribute to unique appearance ideals that further potentiate ED risk among gay men (Murray et al., 2017). The development of specific norms for ED attitudes and behaviours among gay men is an important

endeavor that may better contextualize research in this population. The aim of the present study was to therefore report community norms for the EDE-Q, the field's most widely used ED symptom measure, among cisgender gay men (i.e., gay men who were assigned male sex at birth and have a gender identity of man).

2 | METHODS

2.1 | Study population

The PRIDE Study is a large-scale national longitudinal cohort study of sexual and gender minority adults, which include but are not limited to people who identify as lesbian, gay, bisexual, transgender, and/or queer in the United States. The PRIDE Study launched in 2017, and data were collected on a secure, cloud-based, web-responsive platform accessible from any computer, tablet, or smartphone. PRIDEnet, a national network of organizations and individuals, was created to actively engage sexual and gender minority communities in all stages of research for the PRIDE Study. Participants in The PRIDE Study were recruited through PRIDEnet constituents, digital communications (blog posts, newsletters), distribution of the PRIDE Study-branded promotional items, in-person outreach at conferences and events, social media advertising, and word-of-mouth. Additional details about The PRIDE Study research platform, recruitment, and design have been previously described (Lunn et al., 2019; Lunn et al., 2019). All PRIDE Study participants were invited to complete the “Eating and Body Image” questionnaire from April 2018 to August 2018. For this analysis, we included participants who reported a male sex assigned at birth, exclusively indicated “man” as their gender identity and exclusively indicated “gay” as their sexual orientation (Appendix A). Participants who reported multiple gender identities or sexual orientations were excluded. Of the 10,665 participants in The PRIDE Study at that time, 4,285 completed the questionnaire. Of these, 1,090 identified as cisgender gay men. However, due to the presence of missing values, only data from 978 cisgender gay men were included in The current study. No compensation was received for questionnaire

completion. This study was approved by the University of California, San Francisco and Stanford University School of Medicine Institutional Review Boards as well as The PRIDE Study's Research Advisory Committee and Participant Advisory Committee.

2.2 | Measures

The EDE-Q is a self-report questionnaire that assesses disordered eating attitudes and behaviours over the previous 28 days (Fairburn & Beglin, 2008). The measure provides a global score and four subscale scores: restraint, eating concern, shape concern, and weight concern. Responses are on a 7-point scale; higher scores reflect greater eating-related concerns or behaviours. Frequencies of disordered eating behaviours (e.g., binge eating, compensatory behaviours) are assessed. In this study, Cronbach's alpha was .93 for the global score, .78 for the restraint subscale, .82 for the eating concern subscale, .82 for the weight concern subscale, and .89 for the shape concern subscale.

The frequency of binge eating and compensatory behaviours was assessed in terms of the number episodes occurring during the past 4 weeks (28 days), in accordance with previous literature (Lavender et al., 2010; Penelo, Villarroel, Portell, & Raich, 2012). Any occurrence was defined as ≥ 1 episode in the past 28 days. Regular occurrence of dietary restraint was defined as going for long periods of time (≥ 8 hr) without eating anything to influence shape or weight for ≥ 13 days over the past 28 days (EDE-Q Item 2). Regular occurrence of excessive exercise was defined as exercising in a driven or compulsive way as a means of controlling weight, shape or amount of fat, or burning off calories for ≥ 20 days over the past 28 days. For all other behaviours (objective binge episodes, self-induced vomiting, and laxative misuse), regular occurrence was defined as ≥ 4 occurrences over the past 28 days.

Sociodemographic information (age, race/ethnicity, education), weight, and height were based on self-report. Body mass index (BMI) was calculated using the standard formula weight (kilograms) divided by height (meters) squared ($BMI = \text{weight}/\text{height}^2$). Participants were asked: "Has a mental health professional or physician ever told you that you have an eating disorder such as anorexia nervosa, bulimia nervosa, or binge eating disorder?" If affirmative, participants were asked to specify which type. Options included anorexia nervosa, bulimia nervosa, binge ED, or other/not specified.

2.3 | Data analysis

SPSS 20.0 was used for all analyses and STATA 15.0 was used for figures. Consistent with previous studies

Highlights

- We present norms for the Eating Disorder Examination Questionnaire among cisgender gay men.
- Cisgender gay men scored in the clinically significant range for shape concerns (21%) and weight concerns (11%).
- Participants reported any occurrences of dietary restraint (20%), objective binge episodes (11%), and excessive exercise (10%) in the past 28 days.

(Lavender et al., 2010; Luce, Crowther, & Pole, 2008; Machado, Machado, Gonçalves, & Hoek, 2007), a cut-off score of ≥ 4 as a marker of clinical significance (range 0–6; higher scores indicate greater symptoms) was used for the global score and each of the four EDE-Q subscales. Associations between participant's BMI and EDE-Q (global score and subscale scores) were assessed through the Pearson product-moment correlation coefficient. We calculated norms in a subset ($n = 178$) of young adult cisgender gay men from The PRIDE Study (ages 18–26 years) in order to compare norms with those previously published in young adult men (Lavender et al., 2010). We chose Lavender et al. (2010) as a comparison group as it was the sample of men with published EDE-Q norms that most closely matched the PRIDE Study (i.e., U.S.-based, non-clinical, adult sample). However, Lavender et al. (2010) did not assess sexual orientation; some gay men may have been included. We are unaware of EDE-Q norms published in an exclusively heterosexual community sample of men. Z-tests or Fisher's exact tests were conducted comparing the proportions of individuals who reported each ED behaviour, and independent samples *t* tests were used to compare the global and subscale scores. Locally-weighted scatterplot (lowess) smoothing curves were created to visualize the relationship between age and EDE-Q global and subscale scores given the wide age range of the sample. Two-tailed tests with a *p*-value were set at .05 for significance.

3 | RESULTS

A total of 978 gay cisgender men were included in this study. The median age was 38.9 years (range 18–82). Mean BMI was 27.1 kg/m² ($SD = 6.3$). A total of 80.5% of the participants identified as White, 1.8% as Black/African American, 6.3% as Hispanic/Latino, 3.2%

as Asian or Pacific Islander, 0.6% as Native American/American Indian, and 7.6% as another race or multiracial. In addition, 77.2% of participants had completed a college degree or higher. Overall, 2.9% of participants reported being told by a mental health provider or physician that they had an ED, including anorexia nervosa (1.5%), bulimia nervosa (0.7%), binge ED (0.9%), or other/not specified (0.4%).

Mean scores, standard deviations, and percentile ranks for the EDE-Q subscales and global score are presented in Table 1. Among cisgender gay men of all ages, 5.7% scored in the clinically significant range on the Restraint subscale, 2.1% on the eating concern subscale, 10.5% on

TABLE 1 Distribution of means, standard deviations, and percentile ranks for Eating Disorder Examination Questionnaire global and subscale scores among cisgender gay men from The PRIDE Study ($N = 978$)

	EDE-Q R	EDE-Q EC	EDE-Q WC	EDE-Q SC	EDE-Q global
<i>M</i> (<i>SD</i>)	1.54 (1.43)	0.63 (0.98)	1.91 (1.47)	2.41 (1.62)	1.62 (1.17)
Range	0–6.00	0–6.00	0–6.00	0–6.00	0–5.55
Percentile rank					
5	—	—	-	0.125	0.11
10	—	—	-	0.5	0.26
15	—	—	0.2	0.63	0.39
20	—	—	0.4	0.75	0.51
25	0.2	—	0.6	1	0.63
30	0.4	—	0.8	1.21	0.78
35	0.6	—	1	1.46	0.96
40	0.8	0.2	1.2	1.7	1.13
45	1	0.2	1.4	1.88	1.31
50	1.2	0.2	1.8	2.13	1.45
55	1.4	0.2	2	2.5	1.61
60	1.6	0.4	2.2	2.75	1.78
65	2	0.4	2.6	3	1.94
70	2.4	0.6	2.8	3.38	2.13
75	2.6	0.8	3	3.75	2.34
80	3	1	3.2	4	2.62
85	3.4	1.4	3.6	4.38	2.94
90	3.6	2	4	4.75	3.29
95	4	2.8	4.6	5.25	3.81
99	5.2	4.4	5.6	6	4.81

Abbreviations: EC, eating concern subscale; EDE-Q, Eating Disorder Examination-Questionnaire; EDE-Q global, global score; *M*, mean; R, restraint subscale; SC, shape concern subscale; *SD*, standard deviation; WC, weight concern subscale.

the weight concern subscore, 21.4% on the shape concern subscore, and 4.0% on the global score. BMI was found to be positively associated with EDE-Q scores including restraint ($r = .16$, $p < .001$), eating concern ($r = .28$, $p < .001$), weight concern ($r = .42$, $p < .001$), shape concern ($r = .33$, $p < .001$), and global ($r = .35$, $p < .001$).

Any occurrence (≥ 1 /week) and regular occurrences (≥ 2 /week) of key ED behavioural features and compensatory behaviours among cisgender gay men of all ages are presented in Table 2. Any occurrence of dietary restraint during the past 28 days was observed for almost 20% of the participants, whereas approximately 11% of the participants endorsed any episode of objective binge eating and excessive exercise. Any occurrence of self-induced vomiting (0.6%) and laxative misuse (1.1%) was rarely observed.

Attitudinal subscales and behavioural features of a subsample of young adult cisgender gay men (ages 18–26) from The PRIDE Study sample ($n = 178$) are shown in Table 3. The young adult cisgender gay men in The PRIDE Study scored higher than the Lavender et al. (2010) sample on the weight concern and shape concern subscales as well as the global score. No significant differences were observed between the young adult cisgender gay men subsample of The PRIDE Study and the young adult men in the Lavender et al. (2010) sample for the restraint and eating concern subscales.

There were no differences between the proportion of young adult cisgender gay men in The PRIDE Study and the proportion of the Lavender et al. (2010) sample for dietary restraint behaviors and laxative misuse.

TABLE 2 Proportion of cisgender gay men engaging in disordered eating behaviors among 978 individuals participating in The PRIDE Study

Disordered eating behavior	Any occurrence		Regular occurrence	
	%	<i>n</i>	%	<i>n</i>
Dietary restraint	19.8	194	4.9	48
Objective binge episodes	10.9	107	5.7	56
Self-induced vomiting	0.6	6	0.4	4
Laxative misuse	1.1	11	0.8	7
Excessive exercise	10.1	99	2.0	20

Note. Any occurrence was defined as ≥ 1 episode in the past 28 days. Regular occurrence of dietary restraint was defined as going for long periods of time (≥ 8 hr) without eating anything to influence shape or weight for ≥ 13 days over the past 28 days (EDE-Q Item 2). Regular occurrence of excessive exercise was defined as exercising in a driven or compulsive way as a means of controlling weight, shape or amount of fat, or burning off calories for ≥ 20 days over the past 28 days. For all other behaviours (objective binge episodes, self-induced vomiting, and laxative misuse), regular occurrence was defined as ≥ 4 occurrences over the past 28 days (Lavender et al., 2010; Penelo et al., 2012).

TABLE 3 Comparisons of eating attitudes and disordered eating behaviors in a subsample of cisgender gay men 18–26 years old in The PRIDE Study ($N = 178$) and young men from the Lavender et al. (2010) sample ($N = 404$)

	The PRIDE Study	Lavender et al. (2010)	
Eating attitudes	$M (SD)$		p
EDE-Q global	1.36 (1.06)	1.09 (1.00)	.002
EDE-Q restraint	1.12 (1.30)	1.04 (1.19)	.466
EDE-Q eating concerns	0.51 (0.77)	0.43 (0.77)	.337
EDE-Q weight concerns	1.59 (1.36)	1.29 (1.27)	.010
EDE-Q shape concerns	2.21 (1.53)	1.59 (1.38)	<.001
Disordered eating behaviors	Any occurrence (%)		p
Dietary restraint	21.9	24.0	.581
Objective binge episodes	8.4	25.0	<.001
Self-induced vomiting	0.0	3.2	—
Laxative misuse	1.7	2.7	.566
Excessive exercise	11.2	31.4	<.001

Note. Any occurrence was defined as ≥ 1 episode in the past 28 days (Lavender et al., 2010; Penelo et al., 2012). EDE-Q scores were compared using independent samples t tests. Proportions of disordered eating behaviours were compared with Z tests or Fisher's exact tests.

Abbreviation: EDE-Q, Eating Disorder Examination-Questionnaire.

However, young adult cisgender gay men in The PRIDE Study endorsed significantly lower rates of objective binge episodes and excessive exercise compared with the Lavender et al. (2010) sample. In addition, compared with a

3.2% rate of the Lavender et al. (2010) sample, no participant from the young adult cisgender gay men subsample in the PRIDE Study endorsed self-induced vomiting.


Lowess smoothing curves demonstrate the relationship between age and EDE-Q Global and subscale scores (Figures 1–5). In general, the relationship between age and EDE-Q scores were inverse U shaped, with lowest EDE-Q scores were observed in early young adulthood and late adulthood, with the highest scores in middle adulthood.

4 | DISCUSSION

We summarize community norms for the EDE-Q, a measure of ED attitudes and behaviours, among cisgender gay men. To our knowledge, this is the first study to report community norms of the EDE-Q among cisgender gay men. The few prior studies to report EDE-Q norms in male populations did not assess sexual orientation and were presumably predominantly cisgender heterosexual samples (Hilbert et al., 2012; Lavender et al., 2010; Quick & Byrd-Bredbenner, 2013; Reas et al., 2012).

We found relatively high weight concerns and shape concerns subscale scores among cisgender gay men in the PRIDE Study. These findings are consistent with the growing literature demonstrating body image and ED-related concerns in sexual minority populations (Austin et al., 2009; Calzo, Jerel P. et al., 2018; Diemer, Grant, Munn-Chernoff, Patterson, & Duncan, 2015; Nagata, Garber, Tabler, Murray, & Bibbins-Domingo, 2018; Tabler, Schmitz, Geist, & Nagata, 2019; Watson, Adjei, Saewyc, Homma, & Goodenow, 2017). However, greater

FIGURE 1 Lowess smoothing of age and Eating Disorders Examination Questionnaire (EDE-Q) global score [Colour figure can be viewed at wileyonlinelibrary.com]



endorsement of the attitudinal components of ED symptomatology by gay men did not necessarily equate to greater disordered eating behaviors. EDE-Q norms studies in other populations have demonstrated differences in ED attitudes and behaviours (Lavender et al., 2010; Luce et al., 2008). In addition, some cisgender gay men may have weight and shape concerns related to muscularity and bulking up (Calzo, Corliss, Blood, Field, & Austin, 2013). This may not be reflected in the disordered eating behaviours measured by the EDE-Q, which may be more oriented towards thinness and weight loss. Although a greater constellation of body image concerns was reported in our sample, it is important not to overestimate how this translates to ED behaviors among gay

men. This underscores the importance of sexual orientation-specific norms when interpreting EDE-Q findings in sexual minority populations.

The minority stress theory may explain the high levels of attitudinal ED symptomatology among gay men compared with their heterosexual counterparts. Prejudice and stigma directed towards sexual minorities bring about unique stressors, which can cause adverse health outcomes including mental health disorders (Meyer, 2003). Gay men with EDs have been noted to have high psychiatric comorbidity including depression and anxiety (Feldman & Meyer, 2010; Tabler et al., 2019; Woodside et al., 2001). Disordered eating behaviours may represent deleterious coping with minority stress. Gay men may be




FIGURE 2 Lowess smoothing of age and Eating Disorders Examination Questionnaire (EDE-Q) restraint subscale [Colour figure can be viewed at wileyonlinelibrary.com]




FIGURE 3 Lowess smoothing of age and Eating Disorders Examination Questionnaire (EDE-Q) eating concern subscale [Colour figure can be viewed at wileyonlinelibrary.com]

more likely to view their bodies as sexual objects, to aim to sexually attract others, and therefore may be more vulnerable to experiencing body dissatisfaction compared with heterosexual men (Siever, 1994); this may be a response to minority stress.

We found lower rates of self-induced vomiting and laxative misuse among young cisgender gay men from The PRIDE Study compared with the sample of young men previously reported by Lavender et al. (2010). Some cisgender gay men may have greater desire for muscularity than their heterosexual counterparts (Calzo, J. P. et al., 2013), which may make them less likely to engage in purging behaviours for weight loss. It is important to note that rates of vomiting and laxative misuse in both sam-

ples were relatively low. These differences may also reflect different samples (Lavender et al., 2010 was undergraduate students at a single university) and time periods, as purging behaviours in men may be decreasing over time (Stephen, Rose, Kenney, Rosselli-Navarra, & Weissman, 2014). This contrasts with findings reporting high rates of purging and laxative misuse in gay adolescent boys compared with their heterosexual counterparts (Austin et al., 2009; Austin, Nelson, Birkett, Calzo, & Everett, 2013; Diemer et al., 2015; Watson et al., 2017).

There are limitations to this study. ED assessment in a convenience sample recruited via a web-based platform may limit generalizability but may also enhance responsiveness to a query on a sensitive topic area. Our sample

FIGURE 4 Lowess smoothing of age and Eating Disorders Examination Questionnaire (EDE-Q) weight concern subscale [Colour figure can be viewed at wileyonlinelibrary.com]





FIGURE 5 Lowess smoothing of age and Eating Disorders Examination Questionnaire (EDE-Q) shape concern subscale [Colour figure can be viewed at wileyonlinelibrary.com]



was highly educated, mostly White, and may not be representative of all cisgender gay men in the United States. Selection bias is possible as individuals with more health problems may be more likely to participate in health studies. There are limitations to using the Lavender et al. (2010) sample as a comparison group as these were young adult men in which sexual orientation was not assessed, although we would presume a low overall number of gay men in an undifferentiated sample.

Given that gender identity and sexual orientation disparities are increasingly recognized, establishing normative data for cisgender sexual minority men—such as those who identify as gay—will enable clinicians and researchers to interpret the EDE-Q scores among cisgender gay men. Additional research examining ED-related norms in other sexual minority populations, such as bisexual men, is warranted.

FUNDING

JN is supported by the Pediatric Scientist Development Program (K12 HD000850) supported by the American Academy of Pediatrics and American Pediatric Society and the American Heart Association Career Development (CDA34760281). MC was supported by a Clinical Research Training Fellowship from the American Academy of Neurology and the Tourette Association of America. SBM was supported by the National Institutes of Health (K23 MH115184). EJC was supported by a postdoctoral research fellowship from the Fulbright Commission and the Argentine Ministry of Education. SG is supported by a National Health and Medical Research Council Early Career Fellowship (1121538). JOM was partially supported by K12DK111028 from the National Institute of Diabetes, Digestive, and Kidney Disorders. AF was supported by the National Institute on Drug Abuse (K23DA039800). Research reported in this article was partially funded through a Patient-Centered Outcomes Research Institute (www.pcori.org; PPRN-1501-26848) to MRL. The statements in this article are solely the responsibility of the authors and do not necessarily represent the views of Patient-Centered Outcomes Research Institute, its Board of Governors or Methodology Committee, or the National Institutes of Health.

CONFLICT OF INTEREST

On behalf of all authors, the corresponding author states that there are no conflicts of interest.

ORCID

Jason M. Nagata  <https://orcid.org/0000-0002-6541-0604>

Emilio J. Compte  <https://orcid.org/0000-0002-6803-5950>

Scott Griffiths  <https://orcid.org/0000-0002-6366-3560>

REFERENCES

- Austin, S. B., Nelson, L. A., Birkett, M. A., Calzo, J. P., & Everett, B. (2013). Eating disorder symptoms and obesity at the intersections of gender, ethnicity, and sexual orientation in US high school students. *American Journal of Public Health, 103*(2), 16. <https://doi.org/10.2105/AJPH.2012.301150>
- Austin, S. B., Ziyadeh, N. J., Corliss, H. L., Rosario, M., Wypij, D., Haines, J., ... Field, A. E. (2009). Sexual orientation disparities in purging and binge eating from early to late adolescence. *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine, 45*(3), 238–245. <https://doi.org/10.1016/j.jadohealth.2009.02.001>
- Calzo, J. P., Austin, S. B., & Micali, N. (2018). Sexual orientation disparities in eating disorder symptoms among adolescent boys and girls in the UK. *European Child & Adolescent Psychiatry, 27*(11), 1483–1490. <https://doi.org/10.1007/s00787-018-1145-9>
- Calzo, J. P., Corliss, H. L., Blood, E. A., Field, A. E., & Austin, S. B. (2013). Development of muscularity and weight concerns in heterosexual and sexual minority males. *Health Psychology: Official Journal of the Division of Health Psychology, American Psychological Association, 32*(1), 42–51. <https://doi.org/10.1037/a0028964>
- Carper, T. L. M., Negy, C., & Tantleff-Dunn, S. (2010). Relations among media influence, body image, eating concerns, and sexual orientation in men: A preliminary investigation. *Body Image, 7*(4), 301–309. <https://doi.org/10.1016/j.bodyim.2010.07.002>
- Darcy, A. M., Hardy, K. K., Lock, J., Hill, K. B., & Peebles, R. (2013). The eating disorder examination questionnaire (EDE-Q) among university men and women at different levels of athleticism. *Eating Behaviors, 14*(3), 378–381. <https://doi.org/10.1016/j.eatbeh.2013.04.002>
- Diemer, E. W., Grant, J. D., Munn-Chernoff, M. A., Patterson, D. A., & Duncan, A. E. (2015). Gender identity, sexual orientation, and eating-related pathology in a national sample of college students. *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine, 57*(2), 144–149. <https://doi.org/10.1016/j.jadohealth.2015.03.003>
- Fairburn, C. G., & Beglin, S. (2008). Eating disorder examination questionnaire. In C. G. Fairburn (Ed.), *Cognitive behavior therapy and eating disorders* (pp. 309–313). New York: Guilford Press.
- Feldman, M. B., & Meyer, I. H. (2010). Comorbidity and age of onset of eating disorders in gay men, lesbians, and bisexuals. *Psychiatry Research, 180*(2–3), 126–131. <https://doi.org/10.1016/j.psychres.2009.10.013>
- Hadland, S. E., Austin, S. B., Goodenow, C. S., & Calzo, J. P. (2014). Weight misperception and unhealthy weight control behaviors among sexual minorities in the general adolescent population. *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine, 54*(3), 296–303. <https://doi.org/10.1016/j.jadohealth.2013.08.021>
- Hilbert, A., de Zwaan, M., & Braehler, E. (2012). How frequent are eating disturbances in the population? Norms of the eating disorder examination-questionnaire. *PLoS ONE, 7*(1), e29125. <https://doi.org/10.1371/journal.pone.0029125>

- Lavender, J. M., De Young, K. P., & Anderson, D. A. (2010). Eating disorder examination questionnaire (EDE-Q): Norms for undergraduate men. *Eating Behaviors*, 11(2), 119–121. <https://doi.org/10.1016/j.eatbeh.2009.09.005>
- Luce, K. H., Crowther, J. H., & Pole, M. (2008). Eating disorder examination questionnaire (EDE-Q): Norms for undergraduate women. *The International Journal of Eating Disorders*, 41(3), 273–276. <https://doi.org/10.1002/eat.20504>
- Lunn, M. R., Capriotti, M. R., Flentje, A., Bibbins-Domingo, K., Pletcher, M. J., Triano, A. J., ... Obedin-Maliver, J. (2019). Using mobile technology to engage sexual and gender minorities in clinical research. *PLoS ONE*, 14(5), e0216282. <https://doi.org/10.1371/journal.pone.0216282>
- Lunn, M. R., Lubensky, M., Hunt, C., Flentje, A., Capriotti, M. R., Sooksaman, C., ... Obedin-Maliver, J. (2019). A digital health research platform for community engagement, recruitment, and retention of sexual and gender minority adults in a national longitudinal cohort study—The PRIDE study. *Journal of the American Medical Informatics Association: JAMIA*, 26, 737–748. <https://doi.org/10.1093/jamia/ocz082>
- Machado, P. P. P., Machado, B. C., Gonçalves, S., & Hoek, H. W. (2007). The prevalence of eating disorders not otherwise specified. *The International Journal of Eating Disorders*, 40(3), 212–217. <https://doi.org/10.1002/eat.20358>
- Matthews-Ewald, M. R., Zullig, K. J., & Ward, R. M. (2014). Sexual orientation and disordered eating behaviors among self-identified male and female college students. *Eating Behaviors*, 15(3), 441–444. <https://doi.org/10.1016/j.eatbeh.2014.05.002>
- Meyer, I. H. (2003). Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: Conceptual issues and research evidence. *Psychological Bulletin*, 129(5), 674–697. <https://doi.org/10.1037/0033-2909.129.5.674>
- Mitchison, D., & Mond, J. (2015). Epidemiology of eating disorders, eating disordered behaviour, and body image disturbance in males: A narrative review. *Journal of Eating Disorders*, 3, 2-y. <https://doi.org/10.1186/s40337-015-0058-yeCollection 2015>
- Murray, S. B., Nagata, J. M., Griffiths, S., Calzo, J. P., Brown, T. A., Mitchison, D., ... Mond, J. M. (2017). The enigma of male eating disorders: A critical review and synthesis. *Clinical Psychology Review*, 57, 1–11. doi: S0272-7358(17)30137-X [pii]. <https://doi.org/10.1016/j.cpr.2017.08.001>
- Nagata, J. M., Garber, A. K., Tabler, J., Murray, S. B., & Bibbins-Domingo, K. (2018). Prevalence and correlates of disordered eating behaviors among young adults with overweight or obesity. *Journal of General Internal Medicine*, 33(8), 1337–1343. <https://doi.org/10.1007/s11606-018-4465-z>
- Penelo, E., Villarroel, A. M., Portell, M., & Raich, R. M. (2012). Eating disorder examination questionnaire (EDE-Q): An initial trial in Spanish male undergraduates. *European Journal of Psychological Assessment*, 28(1), 76–83. <https://doi.org/10.1027/1015-5759/a000093>
- Quick, V. M., & Byrd-Bredbenner, C. (2013). Eating disorders examination questionnaire (EDE-Q): Norms for US college students. *Eating and Weight Disorders: EWD*, 18(1), 29–35. <https://doi.org/10.1007/s40519-013-0015-1>
- Reas, D. L., Øverås, M., & Rø, O. (2012). Norms for the eating disorder examination questionnaire (EDE-Q) among high school and university men. *Eating Disorders*, 20(5), 437–443. <https://doi.org/10.1080/10640266.2012.715523>
- Siever, M. D. (1994). Sexual orientation and gender as factors in socioculturally acquired vulnerability to body dissatisfaction and eating disorders. *Journal of Consulting and Clinical Psychology*, 62(2), 252–260. <https://doi.org/10.1037//0022-006x.62.2.252>
- Smith, K. E., Mason, T. B., Murray, S. B., Griffiths, S., Leonard, R. C., Wetterneck, C. T., ... Lavender, J. M. (2017). Male clinical norms and sex differences on the eating disorder inventory (EDI) and eating disorder examination questionnaire (EDE-Q). *The International Journal of Eating Disorders*, 50(7), 769–775. <https://doi.org/10.1002/eat.22716>
- Stephen, E. M., Rose, J. S., Kenney, L., Rosselli-Navarra, F., & Weissman, R. S. (2014). Prevalence and correlates of unhealthy weight control behaviors: Findings from the national longitudinal study of adolescent health. *Journal of Eating Disorders*, 2(1), 16. <https://doi.org/10.1186/2050-2974-2-16>
- Tabler, J. T., Schmitz, R., Geist, C., & Nagata, J. M. (2019). Does it get better? Change in depressive symptoms from late-adolescence to early-adulthood, disordered eating behaviors, and sexual identity. *Journal of Gay & Lesbian Mental Health*, (In Press), 23, 221–243. <https://doi.org/10.1080/19359705.2019.1568944>
- Watson, R. J., Adjei, J., Saewyc, E., Homma, Y., & Goodenow, C. (2017). Trends and disparities in disordered eating among heterosexual and sexual minority adolescents. *The International Journal of Eating Disorders*, 50(1), 22–31. <https://doi.org/10.1002/eat.22576>
- Woodside, D. B., Garfinkel, P. E., Lin, E., Goering, P., Kaplan, A. S., Goldbloom, D. S., & Kennedy, S. H. (2001). Comparisons of men with full or partial eating disorders, men without eating disorders, and women with eating disorders in the community. *The American Journal of Psychiatry*, 158(4), 570–574. <https://doi.org/10.1176/appi.ajp.158.4.570>

How to cite this article: Nagata JM, Capriotti MR, Murray SB, et al. Community norms for the Eating Disorder Examination Questionnaire among cisgender gay men. *Eur Eat Disorders Rev*. 2019;1–10. <https://doi.org/10.1002/erv.2708>

APPENDIX A




FIGURE A1 Flowchart of included study participants